


STDs in Minorities

Sexually Transmitted Disease Surveillance 1998

Division of STD Prevention


Chlamydia - Positivity among women tested in family planning clinics by race and ethnicity: Region X, 1988-1998



Note: Women who met screening criteria were tested. Trends not adjusted for changes in laboratory test method in 1994 and 1998 and associated increases in test sensitivity.


SOURCE: Regional Infertility Prevention Program: Region X Chlamydia Project (Alaska, Idaho, Oregon and Washington)

Gonorrhea - Reported rates for 15-19 year old females by race and ethnicity: United States, 1981-1998




Note: Black, White, and Other are non-Hispanic.

Gonorrhea - Reported rates for 15-19 year old males by race and ethnicity: United States, 1981-1998




Note: Black, White, and Other are non-Hispanic.

Primary and secondary syphilis - Reported rates for 15-19 year old females by race and ethnicity: United States, 1981-1998




Note: Black, White, and Other are non-Hispanic.

Primary and secondary syphilis - Reported rates for 15-19 year old males by race and ethnicity: United States, 1981-1998



Note: Black, White, and Other are non-Hispanic.

Congenital syphilis - Rates for infants <1 year of age by race and ethnicity: United States, 1991-1998



Note: Less than 5% of cases had missing race/ethnicity information and were excluded. Black, White, and Other are non-Hispanic.